

Poster presentation

## **The AP-1 binding sites located in the *pol* gene intragenic regulatory region of HIV-1 are important for virus infectivity**

Nathalie Vandenhoudt<sup>\*1</sup>, Stéphane de Walque<sup>1</sup>, Benoît Van Driessche<sup>1</sup>, Laurence Colin<sup>1</sup>, Valérie Martinelli<sup>1</sup>, Allan Guiguen<sup>1</sup>, Caroline Vanhulle<sup>1</sup>, Arsène Burny<sup>1</sup>, Georges Herbein<sup>2</sup>, Olivier Rohr<sup>3</sup> and Carine Van Lint<sup>1</sup>

Address: <sup>1</sup>Laboratory of Molecular Virology, Institut de Biologie et de Médecine Moléculaires (IBMM), Université Libre de Bruxelles (ULB), 6041 Gosselies, Belgium, <sup>2</sup>Department of Virology, University of Franche-Comte, St-Jacques Hospital, 25030 Besançon, France and <sup>3</sup>INSERM U575, Virology Institute, 67000 Strasbourg, France

\* Corresponding author

from *Frontiers of Retrovirology: Complex retroviruses, retroelements and their hosts*  
Montpellier, France. 21-23 September 2009

Published: 24 September 2009

*Retrovirology* 2009, **6**(Suppl 2):P89 doi:10.1186/1742-4690-6-S2-P89

This abstract is available from: <http://www.retrovirology.com/content/6/S2/P89>

© 2009 Vandenhoudt et al; licensee BioMed Central Ltd.

We have previously identified three AP-1 binding sites in the *pol* gene of human immunodeficiency virus type 1 (HIV-1) and shown that short oligonucleotides containing these sites functioned as phorbol ester-inducible enhancers (Van Lint et al., 1991, *J. Virol.*, 65:7066-7072). These sites are located in a region, called fragment 5103, exhibiting a phorbol ester-inducible enhancing activity on the viral thymidine kinase promoter in HeLa cells. In this study, we have further characterized each of the AP-1 binding sites and have shown that transcription factors c-Fos, JunB and JunD interacted *in vitro* with these motifs. For each site, we have identified mutations abolishing AP-1 factor binding without altering the underlying amino acid sequence of the HIV-1 reverse transcriptase. By transient transfection assays, we have demonstrated that the intragenic AP-1 binding sites were entirely responsible for the PMA-dependent transcriptional activity of fragment 5103. Moreover, this PMA-stimulated activity of fragment 5103 was inhibited by a dominant-negative A-Fos mutant provided the AP-1 sites were not mutated. Finally, we have investigated the biological significance of the intragenic AP-1 binding sites in HIV-1 replication and have shown that these sites are important for viral infectivity.